

Title	Nisin M: a bioengineered Nisin A variant that retains full induction capacity but has significantly reduced antimicrobial activity
Authors	O'Connor, Michelle;Field, Des;Grainger, Aoife;O'Connor, Paula M.;Draper, Lorraine;Ross, R. Paul;Hill, Colin
Publication date	2020-07-20
Original Citation	O'Connor, M., Field, D., Grainger, A., O'Connor, P. M., Draper, L., Ross, R. P. and Hill, C. (2020) 'Nisin M: a bioengineered Nisin A variant that retains full induction capacity but has significantly reduced antimicrobial activity', Applied and Environmental Microbiology, 86(15), e00984-20 (13pp). doi: 10.1128/AEM.00984-20
Type of publication	Article (peer-reviewed)
Link to publisher's version	10.1128/AEM.00984-20
Rights	© 2020, American Society for Microbiology. All Rights Reserved.
Download date	2023-05-04 22:28:26
Item downloaded from	http://hdl.handle.net/10468/12183

1 Nisin M is a bioengineered Nisin A variant that retains full induction capacity but has
2 significantly reduced antimicrobial activity.

3

4 Michelle O' Connor^{1,2†}, Des Field^{1,2†*}, Aoife Grainger¹, Paula M. O' Connor^{2,3}, Lorraine
5 Draper^{1,2}, R. Paul Ross^{1,2}, Colin Hill^{1,2**}

6

7 ¹School of Microbiology, University College Cork, Cork, Ireland

8 ²APC Microbiome Ireland, University College Cork, Cork, Ireland

9 ³Teagasc Food Research Centre, Moorepark, Fermoy, Cork, Ireland

10

11 Address correspondence to:

12 *Des Field, des.field@ucc.ie

13 ** Colin Hill, c.hill@ucc.ie

14 † Michelle O' Connor and Des Field contributed equally to this work. Author order was based on
15 majority of work carried out.

16

17 Keywords:

18 Nisin, signal transduction, induction capacity, antimicrobial activity, NICE system, protein
19 expression

20 Abstract

21 Nisin A is a potent antimicrobial with potential as an alternative to traditional antibiotics, and a
22 number of genetically modified variants have been created that target clinically relevant
23 pathogens. In addition to antimicrobial activity, nisin auto-regulates its own production via a
24 signal transduction pathway, a property that has been exploited in a protein expression system
25 termed the Nisin Controlled Gene Expression (NICE) system. Although NICE has become one
26 of the most popular protein expression systems, one drawback is that the inducer peptide, nisin
27 A, also has inhibitory activity. It has already been demonstrated that the N-terminal region of
28 nisin A contributes to antimicrobial activity and signal transduction properties, therefore, we
29 conducted bioengineering of nisin at positions Pro9 and Gly10 within ring B to produce a bank
30 of variants that could potentially be used as alternative induction peptides. One variant,
31 designated nisin M, has threonines at positions 9 and 10 and retains induction capacity
32 comparable to the wild type nisin A, while most of the antimicrobial activity is abolished.
33 Further analysis confirmed that nisin M produces a mix of peptides as a result of different
34 degrees of dehydration of the two threonines. We show that nisin M exhibits potential as a more
35 suitable alternative to nisin A for the expression of proteins that may be difficult to express, or to
36 produce proteins in strains that are sensitive to wild type nisin. Moreover, it may address the
37 increasing demand by industry for optimization of peptide fermentations to increase yields or
38 their production rate.

39

40

41

42

43 Importance

44 This study describes the generation of a nisin variant with superior characteristics for use in the
45 NICE protein expression system. The variant, termed nisin M, retains an induction capacity
46 comparable to the wild type nisin A but exhibits significantly reduced antimicrobial activity and
47 can therefore be used at concentrations that are normally toxic to the expression host.

48

49

50

51

52

53

54

55

56

57

58

59

60

61

62 Introduction

63 Producing high quantities of proteins of biotechnological and pharmaceutical value from their
64 natural sources can have economic challenges. Although *Escherichia coli* has been the dominant
65 player in the production of recombinant proteins for decades, several issues including the
66 presence of endotoxin or lipopolysaccharide requires expensive and often problematic
67 downstream purification processes (1). The lactic acid bacteria (LAB) *Lactococcus lactis* has
68 gained importance as a host for heterologous protein expression due to its well understood
69 genetics and metabolism, generally regarded as safe (GRAS) status, as well as the availability of
70 a wide range of genetic tools. Indeed, a major advance with regards to protein expression in *L.*
71 *lactis* was the discovery and use of gene expression systems based on a number of inducible
72 promoters. These include promoters that respond to the environment such as P170, which is
73 upregulated at low pH (2) and zinc-based systems that respond to zinc availability (3). One of
74 the best known and most widely employed expression systems is the nisin-inducible controlled
75 gene expression (NICE) system (4, 5) which stems from the nisin biosynthetic operon
76 (*nisABTCIPRKFE*) found in some *L. lactis* strains (6). Nisin is a 34 amino acid peptide and is
77 the most extensively studied bacteriocin (ribosomally synthesized, antimicrobial peptides
78 produced by bacteria) (7, 8). It targets a wide range of Gram-positive bacteria, including food
79 pathogens such as *Listeria monocytogenes*, *Staphylococcus aureus*, *Bacillus cereus* and clostridia
80 (9, 10). Nisin induces its own biosynthesis via a two-component signal transduction pathway
81 NisRK (6) and has led to the development and application of a food grade expression system
82 using *L. lactis* as the host (11). The NICE system encompasses both regulatory elements of the
83 nisin operon, *PnisA*, the nisin-inducible promoter (cloned into several expression vectors) and
84 *nisRK*, the two component histidine kinase response regulator system (harboured by compatible

85 plasmids or inserted on the chromosome of a suitable host strain). The system is ‘switched on’
86 by the addition of nisin in the nanomolar range which activates the receptor NisK. NisK activates
87 NisR by phosphorylation and the activated NisR induces expression at the nisin A promoter (4).
88 The NICE system has been extensively used to produce proteins in *L. lactis*, such as
89 bacteriophage lysins and metalloendopeptidases to demonstrate their potential in dairy
90 fermentations (12, 13). Moreover, NICE can, under certain conditions and with some
91 modifications to the system components, also be used in other species of LAB and in other
92 Gram-positive bacteria (11, 14). Its numerous advantages include ease of use, exquisitely
93 controlled and efficiently induced expression and amenability to large-scale production
94 processes. As an example, nisin induced fermentations of the antimicrobial lysostaphin have
95 been carried out and even identified areas of the NICE system that need improvement (5).
96 However, for industrial applications, nisin addition remains costly (15). Another drawback of the
97 system is that the inducing peptide is also toxic due to its potent antimicrobial activity.
98 Therefore, a nisin peptide that retains its induction capacity whilst having little to no
99 antimicrobial activity would be highly desirable. The gene-encoded nature of the nisin peptide
100 makes genetic engineering to develop certain characteristics of the molecule an attractive and
101 feasible option. Although the bioengineering of nisin commenced over three decades ago, the
102 majority of studies have largely focused on identifying nisin variants with enhanced
103 antimicrobial activity or an extended-antimicrobial spectrum (10, 16, 17). The importance of the
104 N-terminus rings A and B with respect to induction has been highlighted on a number of
105 occasions (6, 16, 18). These studies involved either combinatorial saturation mutagenesis of
106 rings A and B (18) or the application of alanine scanning approaches to assess the antimicrobial
107 activity and induction properties of various nisin derivatives (16).

108 In this study, we carried out a more comprehensive bioengineering approach and created banks
109 of nisin derivatives that have been randomized at positions 9 (P9X) and 10 (G10X) individually
110 and in combination (P9XG10X) and assessed them for antimicrobial activity in conjunction with
111 their ability to induce the nisin promoter using GFP and β -galactosidase reporter systems.

112

113

114

115

116

117

118

119

120

121

122

123

124

125

126

127 Results

128 Creation and screening nisin derivatives for antimicrobial activity and induction capacity.

129 Previous studies utilising site-directed and alanine scanning mutagenesis of nisin have revealed
130 that the N-terminal ring structures are an important region required to activate NisRK (16, 18). In
131 particular, mutagenesis of ring B has been shown to modulate antimicrobial and induction
132 activity. We selected this location as a suitable target for the generation of variants to screen for
133 our desired activities. In order to fully exploit the potential of the nisin ring B we undertook a
134 complete randomisation of the two amino acids in Pro9 and Gly10, both alone (P9X, G10X) and
135 in combination (P9XG10X) using NNK scanning of both codons in the nisin A structural gene
136 (*nisA*) as previously described (7). A bank consisting of 1,452 individual variants created in *L.*
137 *lactis* NZ9800 pCI372*nisA* (pDF05) were screened for antimicrobial activity using deferred
138 antagonism agar diffusion assays and their ability to induce the nisin promoter, *PnisA* fused to a
139 *gfp* reporter gene. The impact of mutations targeting position nine (proline) on antimicrobial
140 activity was assessed using an overlay assay and resulted in zones ranging from those
141 comparable to the wild type control to those devoid of any observable activity. Analysis of
142 colonies using mass spectrometry and/or DNA sequencing identified 12 different amino acid
143 substitutions corresponding to P9H, P9E, P9S, P9T, P9N, P9A, P9M, P9I, P9V P9L, P9W, and
144 P9F (Figure 1A). Substituting P9 with an alanine (P9A) had no impact on either antimicrobial
145 activity or induction capacity. A number of variants (P9H, P9E, P9W and P9F) displayed a loss
146 of both properties. Several others (P9M, P9L, P9N, P9V and P9I) displayed a significant
147 reduction in antimicrobial activity (between 50-65% of wild type) as well as a reduced ability to
148 induce the *gfp* reporter. Notably, P9T and P9S exhibited a slight reduction in antimicrobial

149 activity (70-75%) but retained 100% and 75% induction capacity respectively compared to the
150 wild type control (Figure 1A), which was in agreement with previous studies (16, 18). It is
151 significant that replacement of P9 with either threonine or serine introduces hydroxylated
152 residues which could act as substrates for the lanthionine modification machinery. Indeed,
153 colony mass spectrometry (CMS) of the P9T and P9S producers revealed the presence of masses
154 corresponding to the presence of both unmodified (threonine or serine) and modified residues
155 (dehydrobutyrine (Dhb) or dehydroalanine (Dha)) (Table1).

156 Analysis of 144 variants where position 10 (glycine) was targeted, revealed that the majority of
157 clones exhibited either wild type activity or displayed a complete loss of both antimicrobial
158 activity and induction capacity. Mass Spectrometry (MS) and DNA sequencing determined that
159 almost all of the active variants had retained the original glycine at position 10 (wild type), but
160 we also detected variants corresponding to G10T and G10S (Figure 1B). Here too, CMS
161 identified a mixture of both modified and non-modified residues in the case of G10S (i.e. G10S
162 and G10Dha) but this was not observed when threonine was present at position 10. A variant that
163 displayed little reduction in activity (>50%) had an alanine (G10A) substitution (Figure 1B). A
164 selection of variants that lacked both antimicrobial activity and induction capacity were
165 subjected to DNA sequencing analysis, which identified substitutions corresponding to G10F,
166 G10W and G10L. The inability to detect as wide a range of active variants at this position may
167 arise from the fact that several variants in this position (including G10D, G10N, G10H, G10R,
168 G10L and G10P) have been linked with the loss of threonine dehydration at position 8, meaning
169 that ring B does not undergo cyclization (18).

170 We then set out to vary both residues 9 and 10 simultaneously. As expected, screening of the
171 doubly randomized P9XG10X bank revealed far fewer bioactive variants (approx. 5.6% of the

172 total) of which the majority were wild type (38/64). The remainder exhibited varying degrees of
173 antimicrobial activity ranging from 10-50% with a concomitant loss in induction capacity (data
174 not shown). However, one clone was conspicuous in that despite its apparent lack of
175 antimicrobial activity, it retained an induction capability comparable to the wild type nisin A
176 (Figure 2A). DNA sequencing analysis revealed a variant corresponding to P9T/G10T (both
177 residues replaced with a threonine, Fig 2B). Furthermore, CMS revealed the presence of masses
178 corresponding to the doubly-modified TT peptide (Dhb9Dhb10) but also species with one
179 modified residue to Dhb and a mass close to a peptide with no modified threonine residues
180 (Figure 2A; Table 1). Purification of the derivative P9T/G10T, we termed nisin M, was carried
181 out with our standard nisin purification protocol and subsequent high-performance liquid
182 chromatography (HPLC) evaluation revealed the presence of two major peaks (data not shown).
183 Mass spectrometry analysis of these fractions revealed the presence of one peptide of 3365 Da
184 (consistent with the presence of two Dhb's) and a second peptide of 3383 Da (consistent with a
185 peptide with one threonine and one Dhb). Additionally, a mass in close agreement to a non-
186 modified peptide with threonines in both positions was also observed (data not shown).

187

188 Minimum inhibitory concentration (MIC) of nisin M

189 Following HPLC and freeze-drying of combined fractions to obtain pure peptides, MIC assays
190 were carried out using equimolar concentrations of nisin A and nisin M against a range of Gram
191 positive targets including genera into which the NICE system has been previously introduced
192 (Table 2). The MIC was determined to be the lowest concentration of peptide that resulted in the
193 absence of visible growth of the target strain after 16 hours under the appropriate growth
194 conditions. We established that the MIC of nisin M against a standard laboratory indicator *L.*

195 *lactis* HP was $2.5 \mu\text{g mL}^{-1}$, reflecting a 16-fold increase in MIC compared to wild type nisin A
196 ($0.156 \mu\text{g mL}^{-1}$). Nisin M displayed a similar decrease in potency against the *L. lactis* NZ9000
197 *gfp* reporter strain and its isogenic equivalent *L. lactis* NZ9000 (Table 2). Several lactobacilli
198 have been used as hosts of the NICE system including *Lactobacillus plantarum*, *Lactobacillus*
199 *helveticus* and *Lactobacillus brevis* (14, 19). When *Lb. plantarum* UCC16 and *Lb. brevis* SA-
200 C12 were assessed, an MIC of $>2.5 \mu\text{g mL}^{-1}$ and $1.25 \mu\text{g mL}^{-1}$ was observed, demonstrating a
201 >4 -fold and 16-fold decrease in antimicrobial activity for nisin M respectively in comparison to
202 wild type peptide. (Table 2).

203

204 Induction capacity of nisin A and nisin M at 10 ng mL^{-1}

205 Determination of the induction capacity of nisin A and nisin M at a concentration of 10 ng mL^{-1}
206 was performed using two reporter systems, by way of measurement of GFP and β -galactosidase
207 production. There was no statistical difference in the dynamics of RLU detection when the GFP
208 reporter strain was induced with nisin A and nisin M ($P > .05$) (Figure 3A). Similarly, induction
209 of the *βgal*⁺ reporter strain also revealed no significant difference between nisin A and nisin M
210 at equivalent concentrations (10 ng mL^{-1}) ($P > .05$) (Figure 3B). Moreover, the rate of expression,
211 and therefore the rate of induction was identical for both nisin A and nisin M under the
212 conditions tested in both GFP and β -galactosidase assays.

213

214 Induction capacity and effect on growth of nisin reporter strain at higher induction
215 concentrations

216 Next, we employed 50 ng mL⁻¹, 100 ng mL⁻¹ and 300 ng mL⁻¹ of peptide to determine the effect
217 of higher concentrations of nisin A and M on both protein expression and on growth of the
218 expression host. Fluorescence (RLU) was measured to determine GFP expression and
219 absorbance readings at OD_{595nm} were taken to observe growth of induced strains. For each of
220 these higher concentrations a significant difference between the level of induction by nisin A and
221 nisin M ($P < .0005$) was noted (Fig. 4A-C). Interestingly, the highest RLU reading attained was
222 from cells induced with nisin M at a final concentration of 50 ng mL⁻¹. At this concentration
223 induction continued for the course of the experiment (18 hrs), whereas at 100 ng mL⁻¹ and 300 ng
224 mL⁻¹ induction reached maximum and decreased after 10-12 hours. Notably, there was no delay
225 in the rate of GFP expression following induction with nisin M at 50 ng mL⁻¹ and 100 ng mL⁻¹
226 (Figure 4A-B) compared to 10 ng mL⁻¹ (Figure 3A), where fluorescence intensifies at
227 approximately 6 hours post induction for all experiments; meanwhile, there was a minor delay of
228 30 minutes in expression with induction at 300 ng mL⁻¹ (Figure 4C).

229 When the effects on growth were analysed, no significant impact on the growth of the GFP
230 reporter strain was observed following induction with nisin M at 50 ng mL⁻¹ ($P > .05$), 100 ng
231 mL⁻¹ ($P > .05$) and 300 ng mL⁻¹ ($P > .05$). The OD₅₉₅ of both non-induced cells and cells induced
232 with nisin M increased approximately 5 hours post induction. However, induction with nisin A at
233 the same concentrations resulted in a significant lag-time in growth. An increase in OD₅₉₅ was
234 not observed until 7.5, 8 and 10 hours post induction at 50 ng mL⁻¹ ($P < .05$ for comparison of
235 nisin M to WT, $P < .0005$ for comparison of WT to uninduced samples), 100 ng mL⁻¹ ($P < .0005$)
236 and 300 ng mL⁻¹ ($P < .0005$), respectively. It is worth noting that although growth was observed
237 by samples at these times following induction with WT nisin, there was no fluorescence detected
238 from the same samples until 12, 16 and 18 hours, respectively (Figure 4A-C & 5A-C).

239

240

241 Discussion

242 Any new technological advancements to improve the production of protein biopharmaceuticals
243 and industrial enzymes by microorganisms is highly desirable. Potential methods to optimize the
244 efficiency of an inducible gene expression system may involve adjustment of inducer dosage
245 and/or the timing of inducer addition. The Gram-positive NICE system is somewhat unusual in
246 that the inducer peptide also has the capacity to kill the expression host, and thus induction and
247 killing capacity must be balanced. The generation of a nisin derivative that retained its induction
248 properties but with reduced antimicrobial properties would represent a significant improvement
249 to the NICE system that could be applied to more sensitive strains.

250 Previous work, where the focus has been on the nisin peptide itself, involved randomised
251 mutagenesis of rings A and B (18) and described mutants that retained considerable auto-
252 induction abilities but with lower antimicrobial properties (and vice- versa). Similarly, Ge and
253 co-workers (2016) applied a complete alanine scanning mutagenesis approach and reported that
254 the N-terminal ring structures (ring A and ring B) in nisin were involved in activating NisK to
255 act as an inducing molecule (16). In this study we focused our attention on ring B with a more
256 systematic mutagenesis approach to identify novel derivatives with altered activity/induction
257 properties. This proved to be successful in that we identified a nisin variant that retains induction
258 capacity that is comparable to the wild type peptide but exhibits significantly less antimicrobial
259 activity.

260 Notably, another lantibiotic, subtilin is structurally closely related to nisin and contains the same
261 lanthionine ring structure but does not induce *PnisA*. Indeed, in the study by Steiß, Korn, Kötter
262 and Entian (2015) the failure of subtilin to induce the histidine kinase NisK was shown to mostly
263 depend on the presence of an N-terminal tryptophan, as its replacement with the aliphatic amino
264 acid residues isoleucine, leucine, and valine led to activation of NisK (20). This suggests further
265 bioengineering at position 1 and indeed other amino acid locations in the nisin M background
266 could potentially enhance induction and reduce antimicrobial activity even further.

267 Although this study highlighted ring B of nisin as a critical region in our quest to separate
268 antimicrobial activity from induction/pheromone activity, more residue positions could and
269 should be targeted. Studies with the natural variant nisin Z have revealed that derivatives
270 corresponding to T2S and M17W exhibited an 11-fold and 2-fold increase in induction capacity
271 relative to the parent peptide, respectively, while derivatives S5T and S3T had significantly
272 reduced induction capacity (7).

273 A computational approach evaluating the antimicrobial activity, induction capacity, production
274 levels and immunity/sensor kinase components of natural and bioengineered nisin derivatives
275 could provide a blueprint for the design of more efficient peptide inducers. For example, the
276 ability of nisin P, A and H to activate *PnisA* fused to a *gfp* reporter was assessed and found to
277 differ (21). The promoter was more sensitive to nisin A (1 ng mL^{-1} – $1 \text{ } \mu\text{g mL}^{-1}$) than nisin H (10
278 ng mL^{-1} – $1 \text{ } \mu\text{g mL}^{-1}$) and nisin P (100 ng mL^{-1} – $10 \text{ } \mu\text{g mL}^{-1}$). Higher concentrations of nisin P
279 were required to activate the promotor, but it continued to induce promoter activity at higher
280 concentrations ($10 \text{ } \mu\text{g mL}^{-1}$) whereas nisin A and H were capable of inducing the promoter only
281 up to $1 \text{ } \mu\text{g mL}^{-1}$ concentrations of peptides. The ability to use higher concentrations of nisin P is
282 most likely due to its decreased antimicrobial activity as compared with nisin A and nisin H.

283 While this might advocate for the use of nisin P as an alternative inducer to nisin A, the peptide
284 does not induce at the lower and commonly used inducing concentration (10 ng mL^{-1}). Notably,
285 the nisin M mutant generated in this study induces at both low and high concentrations. While
286 no significant difference in growth of the induced strain compared to the un-induced control was
287 observed, even at the maximum concentration applied (300 ng mL^{-1}), further evaluation with
288 even higher concentrations of nisin M are necessary and with a variety of expression host strains.
289 However, the practicality of using such high concentrations in terms of industrial applications
290 would need to be considered, given that cell free supernatant from a nisin M producer would be
291 the most likely option for induction (rather than expensive purified nisin peptides); though a
292 fermentate analogous to nisaplin (2.5% nisin A) would enable a range of concentrations to be
293 applied irrespective of the sensitivity of host strains (e.g. induction levels above 10 ng mL^{-1} nisin
294 A results in inhibitory effects on the expression strain *L. lactis* NZ9000) (22).

295 Additionally, the natural variant nisin Q also displays similar antimicrobial capabilities to that of
296 nisin A but differs in its ability to induce the *nisA* promoter (23). Directed mutagenesis and
297 analysis of the four amino acids which differ between nisin A and nisin Q (A15V, M21L, H27N,
298 I30V) may help us to more completely understand the pheromone activity of nisin. Remarkably,
299 in the study by Ge and co-workers the derivatives L16D, L16A, L16H, L16V, M21A, M21D,
300 and M21N all exhibited enhanced induction properties when assessed by β -Galactosidase assays,
301 with L16D being particularly notable given it also displays a significant reduction in
302 antimicrobial activity (16). Other regions of nisin subjected to bioengineering approaches and
303 shown to impact on induction activity include the C-terminus and in particular serine and
304 isoleucine at positions 29 and 30, respectively (24). Although the specifics of the interaction
305 between the nisin peptide and NisK have yet to be fully elucidated, a recent study has provided

306 some insight through mutational analysis of NisK. Mutagenesis of conserved residues in the
307 extracellular region of NisK revealed that several hydrophobic residues including two aromatic
308 residues (Tyr113 and Phe133) are crucial for NisK in sensing nisin and regulating nisin
309 biosynthesis (25).

310 Elimination of the antimicrobial activity of nisin is a priority when aiming to improve the nisin
311 peptide in terms of its suitability as a peptide inducer, such as in the NICE system. For example,
312 Reunanen & Saris (2003) developed a method for the quantification of nisin in food samples,
313 through the construction of a non-nisin producing *L. lactis* strain (LAC240), with a plasmid
314 containing a *gfp* gene under the control of the *nisF* promoter and the constituent genes of the
315 nisin two-component regulatory system, *nisRK*. It was reported that upon the addition of nisin
316 peptide concentrations greater than 20 ng mL⁻¹, the LAC240 cells became stressed resulting in a
317 reduction in the quantity of GFP produced and the signal reached the background level when the
318 concentration of nisin was approximately 60 ng mL⁻¹ (26). Moreover, in a study that aimed to
319 improve the response of *L. lactis* to freezing damage through expression of an antifreeze peptide
320 (SF-P), the recombinant strain *L. lactis* NZ3900 SF-P was incubated with different
321 concentrations of nisin (25, 50, or 100 ng mL⁻¹) and at various pH and growth temperature
322 values (27). Notably, maximal expression was observed at 25 ng mL⁻¹, with a much lower level
323 of expression at 50 ng mL⁻¹ and virtually no expression at 100 ng mL⁻¹, most likely due to the
324 inhibitory effects of nisin A, though pH and temperature values were also a factor (27). In
325 another study that sought to optimize the NICE system for the expression of lysostaphin for both
326 laboratory (1 L) and industrial-scale (3000 L) applications and at high cell densities, the authors
327 noted that the addition of too much nisin was detrimental for product formation. Notably, when
328 the culture was induced at higher cell densities, 160 mg L⁻¹ lysostaphin was formed with 20 ng

329 mL⁻¹ nisin and 220 mg L⁻¹ lysostaphin was produced when 40 ng mL⁻¹ nisin was used for
330 induction, indicative of a clear correlation between the cell density at induction and the amount
331 of nisin that is needed for maximal induction (5). While this group reported that maximum
332 protein yield in the NICE system is achieved by induction carried out at a cell density of OD₆₀₀ =
333 5 with a final concentration of 40 ng mL⁻¹ of nisin, we suggest that Nisin M provides for a
334 greater flexibility with respect to inducer concentration by virtue of the attenuated antimicrobial
335 activity of the peptides and the application of high concentrations of inducer peptide is not now a
336 limiting factor.

337 To date, a multitude of peptides, enzymes and vaccines of clinical and biotechnological interest
338 have been overexpressed using nisin, including the anti-bacterial protein lysostaphin (5), a
339 haemagglutinin of the H5N1 influenza virus (28) and Rotavirus VP6 Protein (29), to name but a
340 few. Though several improvements have been made to the NICE system, further improvements
341 are possible. For example, streamlined-genome mutants of *L. lactis* NZ9000 were generated by
342 deletion of four large nonessential DNA regions accounting for 2.83% of the genome and
343 evaluated as microbial cell factories for recombinant protein production. Indeed, following nisin
344 induction, not only was the transcriptional efficiency improved but also the production levels of
345 the expressed reporters were approximately three to fourfold enhanced compared with the wild
346 strain (30). Additionally, expression from the $\Delta lacF$ host-strain *L. lactis* NZ3900 (a strain unable
347 to utilize lactose), enabled food-grade, lactose-based plasmid selection and induction (31) whilst
348 deletion of a specific proteinase gene (NZ9000 $\Delta htrA$) led to increased stability of heterologous-
349 secreted proteins (32).

350 While the aforementioned studies focused on improving the host strain for expression of
351 proteins, this study focuses on potential improvements that can be made to the inducing peptide

352 via mutagenesis of ring B, which has already been reported as playing an important role in
353 induction capacity (16, 18). This study has demonstrated that a nisin A variant with
354 modifications to ring B retained comparable induction capacity to the wild type nisin A peptide
355 yet exhibited less inhibitory effects on the growth of the strain *L. lactis* NZ9000 when applied at
356 concentrations as high as 300 ng mL⁻¹ (0.09 µM). It was also determined this combination has
357 between >4 and >16 fold less activity against various genera and species of bacteria into which
358 the NICE system has been introduced, therefore supporting the claim that nisin M exhibits
359 potential as a suitable alternative to nisin A for use in the NICE system.

360 This study confirms that random mutagenesis experiments continue to be beneficial with a view
361 to enhancing the functional properties of the nisin peptide for specific applications and provide
362 novel nisin variants that exhibit potential for future applications in the pharmaceutical,
363 biotechnological and industrial fields.

364

365

366

367

368

369

370

371

372

373

374

375

376 Materials and Methods:

377 Bacterial strains and plasmids

378 Bacterial strains and plasmids used in this study are listed in Table 3.

379 Creation and analysis of a bank of nisin A ring-B derivatives

380 Mutagenesis of the *nisA* gene was carried out as described previously (7). Briefly, saturation381 mutagenesis was carried out using pDF05 (pCI372-*nisA*) as template and using oligonucleotides

382 as listed in (Table 4) containing an NNK codon in place of each native codon. PCR amplification

383 was performed in a total volume of 50 μ L with 0.5 ng of target DNA (pCI372-*nisA*), 1 unit

384 Phusion High-Fidelity DNA polymerase (Finnzymes, Finland), 1 mM dNTPs and 500 ng each of

385 the appropriate forward and reverse oligonucleotides. The reaction was pre-heated at 98°C for

386 2 min, and then incubated for 29 cycles at 98°C for 30 s, 55°C for 15 s and 72°C for 3 min 30 s,

387 and then finished by incubating at 72°C for 3 min 30 s. Amplified products were treated with

388 DpnI (Stratagene) for 60 min at 37°C to digest template DNA and purified using the QIAquick

389 PCR purification kit. Following transformation of *E. coli* Top 10 cells plasmid DNA was isolated

390 and sequenced using primers pCI372FOR and pCI372REV (Table 4) to verify that mutagenesis

391 had taken place. The purified products were subsequently introduced by electroporation into the

392 strain *L. lactis* NZ9800 which has all the genes necessary for nisin production. Approximately

393 150 transformants were chosen at random for each single position (P9X and G10X) and 1152

394 transformants for the randomised P9XG10X bank. Isolated colonies were inoculated into 96-well
395 plates containing GM17 Cm¹⁰, incubated overnight and stored at -20°C after addition of 80%
396 glycerol. Deferred antagonism assays were performed by replicating strains on GM17 agar plates
397 and allowing them to grow overnight before overlaying with GM17 agar (0.75% w/v agar)
398 seeded with the *L. lactis* HP indicator strain. Induction assays were carried out by replicating
399 strains from each 96 well plate into a fresh 96 well plate containing GM17 broth pre-inoculated
400 with *L. lactis* NZ9000 pNZ8150gfp+, in which GFP acts as a reporter of expression from a nisin
401 inducible promoter (24). Induction of GFP was monitored over 20 hours in terms of relative
402 fluorescence units (RFU) using a TECAN Genios Fluorescence, Absorbance and Luminescence
403 Reader using excitation and emission spectra of 485nm and 535nm, respectively.

404

405 MALDI TOF Mass Spectrometry

406 For Colony Mass Spectrometry (CMS), bacterial colonies of P9X and G10X mutants were
407 collected with sterile plastic loops and mixed with 50 µL of 70% IPA containing 0.1%
408 Trifluoroacetic acid (TFA). The suspension was vortexed, the cells centrifuged in a benchtop
409 centrifuge at 8260 g for 2 min and the supernatant was removed for analysis. For MALDI TOF
410 Mass Spectrometry of nisin M cell free supernatant (CFS) was purified prepared as follows; a
411 1% inoculum of nisin mutant producing strains were grown overnight in 50 mL clarified TY
412 broth and incubated overnight at 30°C. Following incubation cells were centrifuged at 5000rpm
413 for 20 mins at 4°C. Cell free supernatant (CFS) was removed and passed through a 1 g (6 mL)
414 Strata C-18 E column (Phenomenex) pre-equilibrated with 6 mL methanol (Fisher Scientific,
415 UK) and 6 mL HPLC grade H₂O. The column was washed with 12 mL 30% ethanol and nisin
416 eluted using 5 mL 70% isopropanol – 0.1% TFA. Mass Spectrometry in all cases was performed

417 with an Axima TOF² MALDI TOF mass spectrometer (Shimadzu Biotech, Manchester, UK). A
418 0.5 µL aliquot of matrix solution (alpha-cyano-4-hydroxycinnamic acid (CHCA), 10 mg mL⁻¹ in
419 50% acetonitrile-0.1% (v/v) TFA) was placed onto the target and left for 1-2 min before being
420 removed. The residual solution was then air dried and the sample solution (re-suspended
421 lyophilised powder or CMS supernatant) was positioned onto the pre-coated sample spot. Matrix
422 solution (0.5 µL) was added to the sample and allowed to air-dry. The sample was subsequently
423 analysed in a positive-ion linear mode.

424

425 Purification of nisin A and nisin M

426 Purifications of nisin A and variant, nisin M were carried out as per a previously employed (33)
427 with modifications. Briefly, overnight cultures of *L. lactis* NZ9800 pDF05*nisM* (APC3920)
428 and *L. lactis* NZ9700 were inoculated at 0.5% into separate purified tryptone-yeast extract (TY)
429 broth (2 × 900 mL) supplemented with 20% glucose and 20% β-glycerophosphate, and incubated
430 at 30°C overnight. Following incubation, the cultures were centrifuged at 6500g at 4°C for 15
431 min. The supernatant was passed through a column containing ~70g Amberlite XAD-16 beads
432 and subsequently washed with 500 mL of 30% ethanol. The nisin was eluted from the column
433 using 70% isopropanol containing 0.1% trifluoroacetic acid (TFA). Simultaneously, bacterial
434 cell pellets were resuspended in 300 mL 70% isopropanol – 0.1% TFA and stirred at room
435 temperature for 3 h. This cell suspension was then centrifuged at 5000g at 4°C for 10 min and
436 the supernatant was retained. The column eluant was pooled with the post-centrifugation
437 supernatant and isopropanol evaporated using a rotary evaporator (BÜCHI Rotavapor R-205,
438 Switzerland). The pH of the sample was adjusted to pH 4.0 and was subsequently passed through
439 a 10 g (60 mL) Strata C-18 E column (Phenomenex) pre-equilibrated with 60 mL methanol

440 (Fisher Scientific, UK) and 60 mL HPLC grade H₂O. After applying 120 mL 30% ethanol, nisin
441 was eluted from the column using 60 ml 70% isopropanol – 0.1% TFA. For HPLC purification
442 12 mL volumes were concentrated to a volume of 2 mL by rotary evaporation and applied to a
443 Phenomenex C12 reverse-phase (RP-HPLC) column (Jupiter 4 µm proteo 90 Å, 250 mm × 10.0
444 mm, 4 µm) previously equilibrated with 25% acetonitrile-0.1% TFA. Nisin was eluted via a
445 gradient of 25–50% acetonitrile-0.1% TFA that was developed from 10–40 min at a flow rate of
446 3.2 mL min⁻¹. Nisin containing fractions were pooled and acetonitrile removed by rotary
447 evaporation. The purified peptides were lyophilised and stored at -20°C.

448

449 Minimum Inhibitory Concentration (MIC) Assays

450 MIC's were also carried out on strains into which the NICE system was reported to have been
451 introduced including, *Lactobacillus plantarum* (*Lb. plantarum*) and *Lactobacillus brevis* (*Lb.*
452 *brevis*) in order to determine the potential of nisin M as an alternative to nisin A in the NICE
453 system.

454 Minimum inhibitory concentration determinations for strains were carried out in triplicate in 96
455 well microtitre plates (Sarstedt) as described previously (34). Plates were pre-treated with bovine
456 serum albumin (BSA) prior to addition of the peptides. Briefly, to each well of the microtitre
457 plate 200 µL of phosphate buffered saline (PBS) containing 1% (w/v) bovine serum albumin
458 (BSA) was added and incubated at 37°C for 30 min. The wells were washed with 200 µL PBS
459 and allowed to dry. Target strains, *L. lactis* spp. cremoris HP, *L. lactis* NZ9000 pNZ8150*gfp*⁺
460 were grown overnight in M17 broth (Sigma) supplemented with glucose (0.5%) at 30°C. *Lb.*
461 *plantarum* and *Lb. brevis* were grown overnight in MRS broth (Oxoid) at 30°C. Strains were

462 sub-cultured into fresh broth and allowed to grow to an OD₆₀₀ of ~0.5, diluted to a final
463 concentration of 10⁵ cfu mL⁻¹ in a volume of 0.2 mL. Nisin A and nisin M peptides were
464 adjusted to a 750 nM starting concentration and 2-fold serial dilutions of each peptide was added
465 to the target strain. After incubation for 16 h at 30°C the MIC was read as the lowest peptide
466 concentration causing inhibition of visible growth.

467

468 Comparison of nisin A and nisin M induction capacity using beta-galactosidase activity
469 β-galactosidase activity assay was performed as previously employed (35) with modifications.
470 Cultures of *L. lactis* NZ9000 pPTPLβgal⁺ were inoculated in M17 broth (Sigma), supplemented
471 with glucose at 0.5% (GM17) and tetracycline (10 μg mL⁻¹), and incubated at 30°C overnight.
472 Following incubation, a 1% inoculum of each replicate was sub-cultured into fresh GM17
473 medium and incubated at 30°C until an OD₆₀₀ of 0.2-0.3 was reached. Cells were then treated
474 separately with nisin A and nisin M purified peptides to a final concentration of 10 ng mL⁻¹.
475 Every hour 1 mL samples of each test were transferred to an eppendorf and centrifuged at 13,000
476 rpm for 2 minutes (Sorvall Legend Micro 17 centrifuge, Thermo Scientific) to harvest cells.
477 Cells were re-suspended in 1 mL lacZ buffer and 0.5 mL of this was treated with 12.5 μL of
478 0.1% SDS and 25 μL of chloroform and incubated at 30°C for 5 minutes to dissolve cell
479 membranes. Following incubation 100 μL of 2-Nitrophenyl-β-D-galactopyranoside (ONPG) (4
480 mg mL⁻¹) (Sigma-Aldrich) was added to each sample and incubated at 37°C until a yellow colour
481 developed. To stop the reaction samples were treated with 250 μL of a 1 M sodium carbonate
482 solution and centrifuged at 8000rpm for 5 minutes (Thermo Scientific). Absorbance readings of
483 supernatant were read at OD₄₂₀ and OD₅₅₀ (SpectraMax M3 spectrophotometer, Molecular

484 Devices, Sunnyvale, California, USA). Measurement of β -galactosidase activity of samples was
485 calculated as $1000 \times (\text{OD}_{420} - [1.75 \times \text{OD}_{550}]) / (t \times v \times \text{OD}_{600})$ as previously described (16).

486

487 Assessment of purified nisin A and nisin M induction capacity using a green fluorescent protein
488 reporter system.

489 Induction assays were performed previously described (24) with modifications. Briefly, cultures
490 of *L. lactis* NZ9000 pNZ8150*gfp*⁺ were inoculated in M17 broth (Sigma), supplemented with
491 glucose at 0.5% (GM17) with chloramphenicol ($10 \mu\text{g mL}^{-1}$) and incubated at 30°C overnight.
492 Following incubation, a 1% inoculum of each replicate was sub-cultured into fresh GM17
493 medium and incubated at 30°C until an OD_{600} of ~ 0.5 was reached. Cells were then diluted to a
494 final concentration of 10^5 cfu mL^{-1} and treated with nisin A and nisin M at final concentrations of
495 10 ng mL^{-1} , 50 ng mL^{-1} , 100 ng mL^{-1} and 300 ng mL^{-1} . Subsequently, 2 mL was transferred to
496 black, 24 well microtitre plates (PerkinElmer) for induction and 200 μL into a 96 well plate
497 (Sarstedt) for absorbance readings. Fluorescence was detected using a SpectraMax M3
498 spectrophotometer (Molecular Devices, Sunnyvale, California, USA) where excitation and
499 emission parameters were set to 485nm and 528nm respectively for fluorescence, while
500 absorbance readings were taken at OD_{595} using a Multiskan FC microplate photometer v1.01.14
501 (Thermo Scientific, Waltham, Massachusetts, USA). Baseline absorbance of un-cultured GM17
502 was subtracted from the fluorescence and absorbance readings of all test samples using SoftMax
503 Pro v6.3 and SkanIt RE v4.1 software, respectively. Fluorescence was reported as relative light
504 units (RLU) and absorbance as $\text{OD}_{595\text{nm}}$. Tests were carried out in triplicate.

505

506

507

508

509 Statistical analysis

510 Statistical analysis was carried out with SPSS Statistics v2. A test of normality was performed to
511 determine data for each test was normally distributed. For normally distributed data a Repeated
512 Measures ANOVA was performed. For data not normally distributed a Levene's test of
513 homogeneity was performed, where if equal variances were assumed the Repeated Measures
514 ANOVA was carried out; and if equal variances were not assumed the non-parametric Friedman
515 test was performed to determine if differences between the two nisin variants induction capacity,
516 and between the growth of the strains when induced with the peptides at higher concentrations
517 compared to an un-induced control were significant. For ANOVA/Friedman's results with a
518 significant difference between groups ($P < .05$) a post hoc test was performed. Post hoc tests for
519 normally distributed/equal variances assumed samples was the Bonferroni test, and for non-
520 normally distributed/equal variances not assumed samples Dunnett's T3 test was performed. The
521 significance threshold for all ANOVA's and non-parametric tests performed was set at .05.

522

523 Acknowledgements

524 This work was supported by the Irish Government under the National Development Plan,
525 through Science Foundation Ireland Investigator awards 10/IN.1/B3027, SFI/12/RC/2273 and
526 SFI/12/RC/2273 P2. (<http://www.sfi.ie>).

DF would like to acknowledge receipt of a Society for Applied Microbiology
(<http://www.sfam.org.uk>) Students into Work Grant for AG. The funders had no role in study
design, data collection and analysis, decision to publish, or preparation of the manuscript.

References

1. Cano-Garrido O, Rueda FL, Sánchez-García L, Ruiz-Ávila L, Bosser R, Villaverde A, García-Fruitós E. 2014. Expanding the recombinant protein quality in *Lactococcus lactis*. Microb Cell Fact **13**:167.
2. Madsen SM, Arnau J, Vrang A, Givskov M, Israelsen H. 1999. Molecular characterization of the pH-inducible and growth phase-dependent promoter P170 of *Lactococcus lactis*. Mol Microbiol **32**:75-87.
3. Llull D, Poquet I. 2004. New expression system tightly controlled by zinc availability in *Lactococcus lactis*. Appl Environ Microbiol **70**:5398-5406.
4. Mierau I, Kleerebezem M. 2005. 10 years of the nisin-controlled gene expression system (NICE) in *Lactococcus lactis*. Appl Microbiol Biotechnol **68**:705-717.
5. Mierau I, Olieman K, Mond J, Smid EJ. 2005. Optimization of the *Lactococcus lactis* nisin-controlled gene expression system NICE for industrial applications. Microb Cell Fact **4**:16.
6. Kuipers OP, Beerthuizen MM, de Ruyter PG, Luesink EJ, de Vos WM. 1995. Autoregulation of nisin biosynthesis in *Lactococcus lactis* by signal transduction. J Biol Chem **270**:27299-27304.

- 549 7. Field D, Connor PM, Cotter PD, Hill C, Ross RP. 2008. The generation of nisin variants
550 with enhanced activity against specific Gram-positive pathogens. *Mol Microbiol* **69**:218-
551 30.
- 552 8. Wirawan RE, Klesse NA, Jack RW, Tagg JR. 2006. Molecular and Genetic
553 Characterization of a Novel Nisin Variant Produced by *Streptococcus uberis*. *Appl*
554 *Environ Microbiol* **72**:1148-1156.
- 555 9. Naghmouchi K, Drider D, Baah J, Teather R. 2010. Nisin A and Polymyxin B as
556 Synergistic Inhibitors of Gram-positive and Gram-negative Bacteria. *Probiotics*
557 *Antimicrob Proteins* **2**:98-103.
- 558 10. Field D, Cotter PD, Ross RP, Hill C. 2015. Bioengineering of the model lantibiotic nisin.
559 *Bioengineered* **6**:187-92.
- 560 11. De Ruyter P, Kuipers OP, De Vos WM. 1996. Controlled gene expression systems for
561 *Lactococcus lactis* with the food-grade inducer nisin. *Appl Environ Microbiol* **62**:3662-
562 3667.
- 563 12. de Ruyter PG, Kuipers OP, Meijer WC, de Vos WM. 1997. Food-grade controlled lysis
564 of *Lactococcus lactis* for accelerated cheese ripening. *Nat Biotechnol* **15**:976-979.
- 565 13. Hickey RM, Ross RP, Hill C. 2004. Controlled autolysis and enzyme release in a
566 recombinant lactococcal strain expressing the metalloendopeptidase enterolysin A. *Appl*
567 *Environ Microbiol* **70**:1744-1748.
- 568 14. Kleerebezem M, Beerthuyzen MM, Vaughan EE, De Vos WM, Kuipers OP. 1997.
569 Controlled gene expression systems for lactic acid bacteria: transferable nisin-inducible
570 expression cassettes for *Lactococcus*, *Leuconostoc*, and *Lactobacillus* spp. *Appl Environ*
571 *Microbiol* **63**:4581-4584.

- 572 15. Özel B, Şimşek Ö, Akçelik M, Saris PE. 2018. Innovative approaches to nisin
573 production. *Appl Microbiol Biotechnol* **102**:6299-6307.
- 574 16. Ge X, Teng K, Wang J, Zhao F, Wang F, Zhang J, Zhong J. 2016. Ligand determinants
575 of nisin for its induction activity. *J Dairy Sci* **99**:5022-5031.
- 576 17. Zhou L, van Heel AJ, Montalban-Lopez M, Kuipers OP. 2016. Potentiating the activity
577 of nisin against *Escherichia coli*. *Front Cell Dev Biol* **4**:7.
- 578 18. Rink R, Wierenga J, Kuipers A, Kluskens LD, Driessen AJ, Kuipers OP, Moll GN. 2007.
579 Dissection and modulation of the four distinct activities of nisin by mutagenesis of rings
580 A and B and by C-terminal truncation. *Appl Environ Microbiol* **73**:5809-5816.
- 581 19. Pavan S, Hols P, Delcour J, Geoffroy M-C, Grangette C, Kleerebezem M, Mercenier A.
582 2000. Adaptation of the nisin-controlled expression system in *Lactobacillus plantarum*: a
583 tool to study in vivo biological effects. *Appl Environ Microbiol* **66**:4427-4432.
- 584 20. Spieß T, Korn SM, Kötter P, Entian K-D. 2015. Activation of histidine kinase SpaK is
585 mediated by the N-terminal portion of subtilin-like lantibiotics and is independent of lipid
586 II. *Appl Environ Microbiol* **81**:5335-5343.
- 587 21. Garcia-Gutierrez E, O'Connor PM, Saalbach G, Walsh CJ, Hegarty JW, Guinane CM,
588 Mayer MJ, Narbad A, Cotter PD. 2020. First evidence of production of the lantibiotic
589 nisin P. *Sci Rep* **10**:1-15.
- 590 22. Kuipers OP, de Ruyter PG, Kleerebezem M, de Vos WM. 1998. Quorum sensing-
591 controlled gene expression in lactic acid bacteria. *J Biotechnol* **64**:15-21.
- 592 23. Yoneyama F, Fukao M, Zendo T, Nakayama J, Sonomoto K. 2008. Biosynthetic
593 characterization and biochemical features of the third natural nisin variant, nisin Q,
594 produced by *Lactococcus lactis* 61-14. *J Appl Microbiol* **105**:1982-1990.

- 595 24. Field D, Blake T, Mathur H, O'Connor PM, Cotter PD, Paul Ross R, Hill C. 2019.
596 Bioengineering nisin to overcome the nisin resistance protein. *Mol Microbiol* **111**:717-
597 731.
- 598 25. Ge X, Teng K, Wang J, Zhao F, Zhang J, Zhong J. 2017. Identification of key residues in
599 the NisK sensor region for nisin biosynthesis regulation. *Front Microbiol* **8**:106.
- 600 26. Reunanen J, Saris P. 2003. Microplate bioassay for nisin in foods, based on nisin-induced
601 green fluorescent protein fluorescence. *Appl Environ Microbiol* **69**:4214-4218.
- 602 27. Zhang L, Jin Q, Luo J, Wu J, Wang S, Wang Z, Gong S, Zhang W, Lan X. 2018.
603 Intracellular Expression of Antifreeze Peptides in Food Grade *Lactococcus lactis* and
604 Evaluation of Their Cryoprotective Activity. *J Food Sci* **83**:1311-1320.
- 605 28. Szczepankowska AK, Szatraj K, Sałański P, Rózga A, Górecki RK, Bardowski JK. 2017.
606 Recombinant *Lactococcus lactis* expressing Haemagglutinin from a polish avian H5N1
607 isolate and its immunological effect in preliminary animal trials. *BioMed Res Int* 2017.
- 608 29. Esteban LE, Temprana CF, Argüelles M, Glikmann G, Castello AA. 2013. Antigenicity
609 and immunogenicity of rotavirus VP6 protein expressed on the surface of *Lactococcus*
610 *lactis*. *BioMed Res Int* 2013.
- 611 30. Zhu D, Fu Y, Liu F, Xu H, Saris PEJ, Qiao M. 2017. Enhanced heterologous protein
612 productivity by genome reduction in *Lactococcus lactis* NZ9000. *Microb Cell Fact* **16**:1.
- 613 31. Platteeuw C, van Alen-Boerrigter I, van Schalkwijk S, De Vos W. 1996. Food-grade
614 cloning and expression system for *Lactococcus lactis*. *Appl Environ Microbiol* **62**:1008-
615 1013.

- 616 32. Lindholm A, Smeds A, Palva A. 2004. Receptor binding domain of Escherichia coli F18
617 fimbrial adhesin FedF can be both efficiently secreted and surface displayed in a
618 functional form in *Lactococcus lactis*. Appl Environ Microbiol **70**:2061-2071.
- 619 33. Smith MK, Draper LA, Hazelhoff P-J, Cotter PD, Ross RP, Hill C. 2016. A
620 bioengineered nisin derivative, M21A, in combination with food grade additives
621 eradicates biofilms of *Listeria monocytogenes*. Front Microbiol **7**:1939.
- 622 34. Field D, Begley M, O'Connor PM, Daly KM, Hugenholtz F, Cotter PD, Hill C, Ross RP.
623 2012. Bioengineered nisin A derivatives with enhanced activity against both Gram
624 positive and Gram negative pathogens. PloS One **7**:e46884.
- 625 35. Israelsen H, Madsen SM, Vrang A, Hansen EB, Johansen E. 1995. Cloning and partial
626 characterization of regulated promoters from *Lactococcus lactis* Tn917-lacZ integrants
627 with the new promoter probe vector, pAK80. Appl Environ Microbiol **61**:2540-2547.
- 628 36. Kuipers OP, Beerthuyzen MM, Siezen RJ, De Vos WM. 1993. Characterization of the
629 nisin gene cluster *nisABTCIPR* of *Lactococcus lactis*: Requirement of expression of the
630 *nisA* and *nisI* genes for development of immunity. Eur J Biochem **216**:281-291.
- 631 37. Åvall-Jääskeläinen S, Kylä-Nikkilä K, Kahala M, Miikkulainen-Lahti T, Palva A. 2002.
632 Surface display of foreign epitopes on the *Lactobacillus brevis* S-layer. Appl Environ
633 Microbiol **68**:5943-5951.
- 634
- 635
- 636
- 637

638 Table 1. Mass spectrometry analysis of selected derivatives

Ring B Derivative	Predicted Mass (Da)	Actual Mass (Da)	Dehydrations		Ref
			Observed	Lacking	
P9A	3328	3327.87	8	0	This study (16)
P9T	3357	3356.67	8	1	This study
		3339.62	9	0	
P9S	3343	3342.73	8	1	This study
		3324.69			
G10A	3366	3367.14	8	0	This study (18)
G10T	3398	3397.76	8	1	This study (18)
G10S	3384	3384.57	8	1	This study (18)
		3367.23	9	0	
P9T/G10T (nisin M)	3402	3399.86	8	2	This study
		3382.91	9	1	
		3365.33	10	0	
Nisin A	3354	3353.44	8	0	

639

640

641 Table 2: MIC of nisin A and nisin M against standard indicator strains (including those reported
642 to have had the NICE system introduced).

Indicator organism	Nisin A $\mu\text{g mL}^{-1}$ (μM)	Nisin M $\mu\text{g mL}^{-1}$ (μM)	Fold decrease in activity
<i>Lb. plantarum</i> UCC16	0.625 (0.1875)	>2.5 (>0.750)	>4
<i>L. lactis</i> NZ9000 pNZ8150gfp+	0.156 (0.046)	>2.5 (>0.750)	>16

<i>L. lactis</i> NZ9000	0.156 (0.046)	2.5 (0.750)	16
pNZ8150			
<i>Lb. brevis</i> SA-C12	0.078 (0.0234)	1.25 (0.375)	16
<i>L. lactis</i> spp. <i>cremoris</i> HP	0.156 (0.0468)	2.5 (0.750)	16

643

644

645 Table 3: Bacterial strains and plasmids used in this study

Strain or Plasmid	Characteristic	Reference
<i>L. lactis</i> NZ9000	MG1363 derivative, <i>NisRK</i> integrated into <i>pepN</i> gene (<i>pepN</i> -). Most commonly used host of the NICE system.	(22) (4)
<i>L. lactis</i> NZ9000 pNZ8150	NZ9000 strain harbouring pNZ8150. pNZ8150: <i>ScaI</i> site for translational fusions, standard vector for NICE system, Cm ^R .	(4)
<i>L. lactis</i> NZ9000 pNZ8150 <i>gfp</i> ⁺	NZ9000 strain harbouring pNZ8150 <i>gfp</i> ⁺ under <i>PnisA</i> promoter. Cm ^R .	(24)
<i>L. lactis</i> NZ9000 pPTPL <i>βgal</i> ⁺	NZ9000 strain harbouring low copy plasmid pPTPL with β -galactosidase expressing gene under the control of the <i>PnisA</i> promoter. Tet ^R	(7)
<i>L. lactis</i> NZ9800	Derivative of NZ9700 with 4bp deletion rendering an inactive nisin operon (Δ <i>nisA</i>), except <i>nisRK</i> genes. Host of the NICE system.	(22, 36)
<i>L. lactis</i> NZ9800 pDF05	NZ9800 harbouring pDF05 (pCI372 with <i>nisA</i> under its own promoter). Wild type nisin A producer, Cm ^R .	(22, 36) (7)

<i>L. lactis</i> NZ9800 pDF05 <i>nisM</i>	pDF05 where codons 9 and 10 of <i>nisA</i> have been randomized. Nisin M producer, Cm ^R .	This work UCC Culture Collection (APC3920)
<i>Lb. plantarum</i> UCC16	Nisin sensitive indicator Species in which NICE system has been utilized.	UCC Culture Collection (4, 19)
<i>Lb. brevis</i> SA-C12	Nisin sensitive indicator Species in which NICE system has been utilized.	UCC Culture Collection (4, 37)
<i>L. lactis</i> ssp. <i>cremoris</i> HP	Nisin sensitive indicator strain	UCC Culture Collection

646

647

648 Table 4. Oligonucleotides utilised in this study.

Primer name	Sequence
NisP9degFOR	5' CTA TGT ACA NNK GGT TGT AAA ACA GGA GCT CTG ATG GGT 3'
NisP9degREV	5' TTT ACA ACC MNN TGT ACA TAG CGA AAT ACT TGT AAT GCG 3'
NisG10degFOR	5' TGT ACA CCC NNK TGT AAA ACA GGA GCT CTG ATG GGT TGT 3'
NisG10degREV	5' TGT TTT ACA MNN GGG TGT ACA TAG CGA AAT ACT TGT AAT 3'
NisP9G10degFOR	5' CTA TGT ACA NNK NNK TGT AAA ACA GGA GCT CTG ATG GGT 3'
NisP9G10degREV	5' TTT ACA MNN MNN TGT ACA TAG CGA AAT ACT TGT AAT

649

650

651

652

653

654

655

656

657

658

659

660

661

662

663

664

665

666

667

668

669

670

671

672

673

674

675

676

677

678

679

680

681

682

683

684

685

686

687

688

689

690

691

692

693

694

695

696

697

698

699

700

701

702

703

704

705

706

707

708

709

710

711

712

713

714

715

716

717

718

719

720

721

722

723

724

725

726

727

728

729

730

731

732

733

734

735

736

737

738

739

740

741

742

743

744

745

746

747

748

749

750

751

752

753

754

755

756

757

758

759

760

761

762

763

764

765

766

767

768

769

770

771

772

773

774

775

776

777

778

779

780

781

782

783

784

785

786

787

788

789

790

791

792

793

794

795

796

797

798

799

800

801

802

803

804

805

806

807

808

809

810

811

812

813

814

815

816

817

818

819

820

821

822

823

824

825

826

827

828

829

830

831

832

833

834

835

836

837

838

839

840

841

842

843

844

845

846

847

848

849

850

851

852

853

854

855

856

857

858

859

860

861

862

863

864

865

866

867

868

869

870

871

872

873

874

875

876

877

878

879

880

881

882

883

884

885

886

887

888

889

890

891

892

893

894

895

896

897

898

899

900

901

902

903

904

905

906

907

908

909

910

911

912

913

914

915

916

917

918

919

920

921

922

923

924

925

926

927

928

929

930

931

932

933

934

935

936

937

938

939

940

941

942

943

944

945

946

947

948

949

950

951

952

953

954

955

956

957

958

959

960

961

962

963

964

965

966

967

968

969

970

971

972

973

974

975

976

977

978

979

980

981

982

983

984

985

986

987

988

989

990

991

992

993

994

995

996

997

998

999

1000

1001

1002

1003

1004

1005

1006

1007

1008

1009

1010

1011

1012

1013

1014

1015

1016

1017

1018

1019

1020

1021

1022

1023

1024

1025

1026

1027

1028

1029

1030

1031

1032

1033

1034

1035

1036

1037

1038

1039

1040

1041

1042

1043

1044

1045

1046

1047

1048

1049

1050

1051

1052

1053

1054

1055

1056

1057

1058

1059

1060

1061

1062

1063

1064

1065

1066

1067

1068

1069

1070

1071

1072

1073

1074

1075

1076

1077

1078

1079

1080

1081

1082

1083

1084

1085

1086

1087

1088

1089

1090

1091

1092

1093

1094

1095

1096

1097

1098

1099

1100

1101

1102

1103

1104

1105

1106

1107

1108

1109

1110

1111

1112

1113

1114

1115

1116

1117

1118

1119

1120

1121

1122

1123

1124

1125

1126

1127

1128

1129

1130

1131

1132

1133

1134

1135

1136

1137

1138

1139

1140

1141

1142

1143

1144

1145

1146

1147

1148

1149

1150

1151

1152

1153

1154

1155

1156

1157

1158

1159

1160

1161

1162

1163

1164

1165

1166

1167

1168

1169

1170

1171

1172

1173

1174

1175

1176

1177

1178

1179

1180

1181

1182

1183

1184

1185

1186

1187

1188

1189

1190

1191

1192

1193

1194

1195

1196

1197

1198

1199

1200

1201

1202

1203

1204

1205

1206

1207

1208

1209

1210

1211

1212

1213

1214

1215

1216

1217

1218

1219

1220

1221

1222

1223

1224

1225

1226

1227

1228

1229

1230

1231

1232

1233

1234

1235

1236

1237

1238

1239

1240

1241

1242

1243

1244

1245

1246

1247

1248

1249

1250

1251

1252

1253

1254

1255

1256

1257

1258

1259

1260

1261

1262

1263

1264

1265

1266

1267

1268

1269

1270

1271

1272

1273

1274

1275

1276

1277

1278

1279

1280

1281

1282

1283

1284

1285

1286

1287

1288

1289

1290

1291

1292

1293

1294

1295

1296

1297

1298

1299

1300

1301

1302

1303

1304

1305

1306

1307

1308

1309

1310

1311

1312

1313

1314

1315

1316

1317

1318

1319

1320

1321

1322

1323

1324

1325

1326

1327

1328

1329

1330

1331

1332

1333

1334

1335

1336

1337

1338

1339

1340

1341

1342

1343

1344

1345

1346

1347

1348

1349

1350

1351

1352

1353

1354

1355

1356

1357

1358

1359

1360

1361

1362

1363

1364

1365

1366

1367

1368

1369

1370

1371

1372

1373

1374

1375

1376

1377

1378

1379

1380

1381

1382

1383

1384

1385

1386

1387

1388

1389

1390

1391

1392

1393

1394

1395

1396

1397

1398

1399

1400

1401

1402

1403

1404

1405

1406

1407

1408

1409

1410

1411

1412

1413

1414

1415

1416

1417

1418

1419

1420

1421

1422

1423

1424

1425

1426

1427

1428

1429

1430

1431

1432

1433

1434

1435

1436

1437

1438

1439

1440

1441

1442

1443

1444

1445

1446

1447

1448

1449

1450

1451

1452

1453

1454

1455

1456

1457

1458

1459

1460

1461

1462

1463

1464

1465

1466

1467

1468

1469

1470

1471

1472

1473

1474

1475

1476

1477

1478

1479

1480

1481

1482

1483

1484

1485

1486

1487

1488

1489

1490

1491

1492

1493

1494

1495

1496

1497

1498

1499

1500

1501

1502

1503

1504

1505

1506

1507

1508

1509

1510

1511

1512

1513

1514

1515

1516

1517

1518

1519

1520

1521

1522

1523

1524

1525

1526

1527

1528

1529

1530

1531

1532

1533

1534

1535

1536

1537

1538

1539

1540

1541

1542

1543

1544

1545

1546

1547

1548

1549

1550

1551

1552

1553

1554

1555

1556

1557

1558

1559

1560

1561

1562

1563

1564

1565

1566

1567

1568

1569

1570

1571

1572

1573

1574

1575

1576

1577

1578

1579

1580

1581

1582

1583

1584

1585

1586

1587

1588

1589

1590

1591

1592

1593

1594

1595

1596

1597

1598

1599

1600

1601

1602

1603

1604

1605

1606

1607

1608

1609

1610

1611

1612

1613

1614

1615

1616

1617

1618

1619

1620

1621

1622

1623

1624

1625

1626

1627

1628

1629

1630

1631

1632

1633

1634

1635

1636

1637

1638

1639

1640

1641

1642

1643

1644

1645

1646

1647

1648

1649

1650

1651

1652

1653

1654

1655

1656

1657

1658

1659

1660

1661

1662

1663

1664

1665

1666

1667

1668

1669

1670

1671

1672

1673

1674

1675

1676

1677

1678

1679

1680

1681

1682

1683

1684

1685

1686

1687

1688

1689

1690

1691

1692

1693

1694

1695

1696

1697

1698

1699

1700

1701

1702

1703

1704

1705

1706

1707

1708

1709

1710

1711

1712

1713

1714

1715

1716

1717

1718

1719

1720

1721

1722

1723

1724

1725

1726

1727

1728

1729

1730

1731

1732

1733

1734

1735

1736

1737

1738

1739

1740

1741

1742

1743

1744

1745

1746

1747

1748

1749

1750

1751

1752

1753

1754

1755

1756

1757

1758

1759

1760

1761

1762

1763

1764

1765

1766

1767

1768

1769

1770

1771

1772

1773

1774

1775

1776

1777

1778

1779

1780

1781

1782

1783

1784

1785

1786

1787

1788

1789

1790

1791

1792

1793

1794

1795

1796

1797

1798

1799

1800

1801

1802

1803

1804

1805

1806

1807

1808

1809

1810

1811

1812

1813

1814

1815

1816

1817

1818

1819

1820

1821

1822

1823

1824

1825

1826

1827

1828

1829

1830

1831

1832

1833

1834

1835

1836

1837

1838

1839

1840

1841

1842

1843

1844

1845

1846

1847

1848

1849

1850

1851

1852

1853

1854

1855

1856

1857

1858

1859

1860

1861

1862

1863

1864

1865

1866

1867

1868

1869

1870

1871

1872

1873

1874

1875

1876

1877

1878

1879

1880

1881

1882

1883

1884

1885

1886

1887

1888

1889

1890

1891

1892

1893

1894

1895

1896

1897

1898

1899

1900

1901

1902

1903

1904

1905

1906

1907

1908

1909

1910

1911

1912

1913

1914

1915

1916

1917

1918

1919

1920

1921

1922

1923

1924

1925

1926

1927

1928

1929

1930

1931

1932

1933

1934

1935

1936

1937

1938

1939

1940

1941

1942

1943

1944

1945

1946

1947

1948

1949

1950

1951

1952

1953

1954

1955

1956

1957

1958

1959

1960

1961

1962

1963

1964

1965

1966

1967

1968

1969

1970

1971

1972

1973

1974

1975

1976

1977

1978

1979

1980

1981

1982

1983

1984

1985

1986

1987

1988

1989

1990

1991

1992

1993

1994

1995

1996

1997

1998

1999

2000

2001

2002

2003

2004

2005

2006

2007

2008

2009

2010

2011

2012

2013

2014

2015

2016

2017

2018

2019

2020

2021

2022

2023

2024

2025

2026

2027

2028

2029

2030

2031

2032

2033

2034

2035

2036

2037

2038

<

665 shown as percentages (%) and ordered from highest to lowest based on biological activity of the
666 variants.

667

668 Figure 2. A: (Top) Biological activity of nisin A and nisin M as determined by deferred
669 antagonism assays and assessment of induction capacity following induction of a *L. lactis* strain
670 containing *gfp*⁺ under control of the nisin promoter. (Bottom) Colony Mass Spectrometry of the
671 wild type nisin A producer (3353.44 Da) and nisin M comprising of a combination of
672 unmodified peptide, single dehydration or two dehydrations at P9T/G10T (3399.86 Da, 3382.91
673 Da, and 3365.33 Da respectively). B: Structure of nisin A where amino acids are represented by
674 their single letter codes and modified residues are indicated as follows; Dha: dehydroalanine,
675 Dhb: dehydrobutyrine, Abu: 2-aminoabutyric acid, Ala-S-Ala: lanthionine, Abu-S-Ala:
676 methylanthionine. Residues in orange and pink show amino acid substitutions for nisin M,
677 producing 4 possible forms of the peptide.

678

679 Figure 3. Induction capacity of nisin A (red/circle) and nisin M (blue/square) determined by
680 expression of (A) GFP and (B) β -galactosidase reporter genes under the control of the *PnisA*
681 promoter in *L. lactis* NZ9000 pNZ8150*gfp*⁺ and *L. lactis* NZ9000 pPTPL *β gal*⁺ respectively
682 when induced at a final concentration of 10 ng mL⁻¹. Negative controls (green/triangle) are
683 uninduced test strains. Statistical analysis shows there is no significant difference between the
684 induction capacities of nisin M and nisin A in both methods tested ($P > .05$).

685

686 Figure 4. Comparison of induction capacities of nisin A (red/circle), nisin M (blue/square)
687 determined by expression of GFP under the control of the *PnisA* promoter in *L. lactis*
688 pNZ8150*gfp*⁺ induced at final concentrations of (A) 50 ng mL⁻¹, (B) 100 ng mL⁻¹ and (C) 300
689 ng mL⁻¹. Statistical analysis demonstrates a significant difference between induction capacity of
690 the two peptides at all concentrations tested, ($P < .0005$) Negative control in this assay is
691 uninduced *L. lactis* NZ9000 pNZ8150*gfp*⁺ (green/triangle).

692

693 Figure 5. Effects of nisin A (red/circle) and nisin M (blue/square) on growth of *L. lactis* NZ9000
694 pNZ8150*gfp*⁺ induced at concentrations of (A) 50 ng mL⁻¹, (B) 100 ng mL⁻¹ and (C) 300 ng mL⁻¹
695 compared to an un-induced control (green/triangle) determined by absorbance at OD_{595nm}.
696 Results show no significant difference between growth of the uninduced control and cells
697 induced with nisin M at all concentrations tested ($P > .05$); while there is a significant difference
698 between the growth of cells induced with WT nisin compared to both the uninduced control (50
699 ng mL⁻¹: $P < .0005$; 100 ng mL⁻¹: $P < .0005$; 300 ng mL⁻¹: $P < .0005$), and samples induced with
700 nisin M (50 ng mL⁻¹: $P < .05$; 100 ng mL⁻¹: $P < .0005$; 300 ng mL⁻¹: $P < .0005$).


701

702

703

704

A



B

